

A Study on Bio-efficacy of eco-friendly pesticides against major predators of lac insect

Shantha A R

Department of Zoology, Sahyadri Science College, Shivamogga, Karnataka, India

To Cite this Article

Shantha A R, "A Study on Bio-efficacy of eco-friendly pesticides against major predators of lac insect", *International Journal for Modern Trends in Science and Technology*, Vol. 05, Issue 10, October 2019, pp.-60-63.

Article Info

Received on 25-September-2019, Revised on 18-October-2019, Accepted on 25-October-2019, Published on 31-October-2019.

ABSTRACT

Lac culture is the most efficient system because it gives jobs to the farmers, it gives pesticides and fertilizers and require less amount of water and resin produced by lac is economically useful. It is used in medicine, ayurvedic and in many fields. They do not harm the environment and other plants. It has much demand in many countries. The insecticidal sprays were applied at 45 days after the inoculation of brood lac on the completion of male emergence. The first spray application was applied at 7 days after the completion of male emergence followed by a second application at 30 days after the first spray. The insecticidal application was applied on lac bearing Twigs of the ber plants to cover the encrustation of lac insect. Among the insecticides evaluated, the most effective treatment was spinosad 2.5% EC against Predators viz., *Eublemma amabilis* and *Pseudohypatopa pulverea* with 84.85 and 78.95 mean per cent reduction in the population over control, respectively. Application of spinosad 2.5% EC was also recorded as the most effective treatment in terms of survival percentage of lac insect. Emamectin benzoate 5% SG was found next effective treatment, whereas, Neem oil was found least effective treatment in reducing the population of predators of lac insect as well as in terms of survival percentage of lac insect.

KEYWORDS: *Eublemma amabilis*, *Pseudohypatopa pulverea*, *Kerria lacca*

Copyright © 2019 International Journal for Modern Trends in Science and Technology
All rights reserved.

I. INTRODUCTION

The versatility and environmental benefits of lac make it a valuable natural resource that has contributed significantly to human welfare throughout history. The continued cultivation and utilization of lac reflect the importance of understanding and harnessing the potential of natural products provided by various living organisms, including insects, for the betterment of society

The lac insect *Kerria lacca* is scale insect which occurs naturally as parasites on various host

plants. The specific host trees of lac insects are palas, ber, kusum, semialata. It secretes lac, a layer of red resin on branches of host-trees on which it settles. Lac resin is a natural, biodegradable, non-toxic, and the only animal originates resin which is used as food, textiles industries and pharmaceutical industries. It is also used as surface coating, electrical component manufacturing, and other fields. The distribution range of lac is tropical and subtropical areas of south and south-east Asia. Lac culture is useful as resin, dye, wax and having tremendous export potential. In India, two species of lac strains

namely Rangeeni and Kusumi are known to prevail dominantly. Of the two strains i.e., Kusumi and Rangeeni of lac insect, Aghani crop of Kusmi contribute the most with the contribution of 32 % followed by Jethwi (26 %) of Kusmi strain and Baisakhi (24 %) and Katki (18 %) of Rangeeni strain in total lac production [1]. It is a highly remunerative crop, paying high economic returns to the farmers and also foreign exchange to the country through its export. In spite of the wide distribution of the lac insect through the country on different hosts, lac production is limited in the country because of many biotic and abiotic factors; Among biotic factors are predators and parasitoids, while adverse climatic factors create abiotic stresses causing hinderance in lac production. Among the natural enemies Eublemma amabilis Moore (Lepidoptera; Noctuidae), Psuedohypatopa pulverea Meyr (Lepidoptera; Blastobesidae), Chrysopa lacciperda Kimmins and Chrysopa madestes Banks (Chrysopidae; Neuroptera) are the major predators of regular occurrence causing severe losses to lac production [2]. Predators have been estimated to cause around 35 to 40 per cent loss of lac production [3, 4], and these are in regular occurrence but their incidence may vary from season to season, place to place and crop to crop. The first, second and third instar larvae of Chrysopa madestes can consume 20, 24 and 74 mature females of lac insect per day, respectively [5]. As many as 30 different species of parasites have been reported on lac insect [6]. Earlier studies indicated a parasitisation level of only 5-10 percent [7], but with changing times and climatic scenario an increased parasitization level of about 20-37 percent has been reported [8], which have been noticed as one of the major limiting factors in complete failure of crop during last few years. Rangeeni crop is more vulnerable to pest attack and the damage is more in the rainy season crop which sometimes destroys the whole crop. In Rajasthan, lac insect has been noticed to prevail naturally on as many as in different hosts [9]. In spite of its high natural occurrence and wide availability of hosts, the lac cultivation is not in practice and yet not adopted by the farmers of the region. Among various factors, the lack of knowledge about practices of lac cultivation, the incidence and management of major predators of lac insect is one of important the factor responsible for the hindering lac production in the region.

Lac cultivation is an important source of livelihood for rural communities, as the lac insects and their products have a wide range of

commercial applications. The lac resin, known as shellac, seedlac, or button lac, finds use in diverse industries such as paints, inks, pharmaceuticals, cosmetics, electrical equipment, automobiles, defence, railways, marine applications, surface coatings, confectionery, and textile dyeing.

Beyond its economic value, lac cultivation is also environmentally sustainable, as it does not require significant inputs of water, pesticides, or fertilizers for the host plants to thrive. The lac insects are considered eco-friendly and play a role in maintaining the ecological balance.

II. MATERIALS AND METHODS

SYSTEMATIC POSITION OF *LACCIFER LACCA*

A number of species of lac insects are known, of this *Laccifer lacca* is by far the most important and produces the bulk of the lac for commerce. It belongs to:

- Phylum – Arthropoda
- Class – Insecta
- Order – Hemiptera
- Super family – Coccoidea
- Family – Lacciferidae
- Genus – *Laccifer*
- Species – *lacca*

The field experiment was conducted to evaluate the bio-efficacy of eco-friendly pesticides against major predators of lac insect and their safety response for the lac insect on ber plants. The ber plants were inoculated with locally collected brood lac after the appearance of new shoots of ber. The brood lac having fully mature female cells on the appearance of the yellow spot was tied on the ber plants for the emergence and settlement of lac insects. There were seven different treatments that were applied to evaluate the response of different botanicals and insecticides against the major predators of lac insect. The insecticidal sprays were applied at 45 days after the inoculation of brood lac on the completion of male emergence. The first spray application was applied at 7 days after the completion of male emergence followed by a second application at 30 days after the first spray. The insecticidal application was applied on lac bearing twigs of the ber plants to cover the encrustation of lac insect. The treatment details are given in table 1. The observation on the efficacy of different treatments were recorded in terms of living/dead cells at harvest. The mature lac stick samples for each treatment in each replication were harvested and observed to count the numbers of mature live

and dead cells per 4 square centimeters by placing the graph paper. The samples were collected from the upper, middle and lower portion of the treated plants for each treatment and kept in 60 mesh nylon cages for the emergence of predators. The numbers of living and dead larvae as well as adults of lac predators emerging from the caged samples were counted, and the percentage reduction in the incidence of predators over control was worked out.

III. RESULT AND DISCUSSION

The bio-efficacy of different insecticides and botanical extracts was evaluated against major natural enemies of lac insect. The first spray was done at 45 days after brood lac inoculation and the subsequent spray was done at 30 days after first spray. The observation on the efficacy of the eco-friendly pesticide on major predators of lac insect was recorded by counting the number of individuals that emerged from the caged samples from each treatment, collected at harvest. All the treatments were significantly superior over control in reducing the population of the major predators of lac insect viz., *Eublemma amabilis* and *Pseudohypatopa pulvereana*.

The results presented in table 2 revealed that the spray application of spinosad 2.5% EC @ 2.0 ml/litre of water proved to be most effective with maximum per cent reduction of the mean population of predators i.e. *E. amabilis* (84.85%) and *P. pulvereana* (78.95) over control with the minimum mean population of 3.33 and 1.33 per meter lac stick. Emamectin benzoate 5% SG and @ 0.4 gm/lit, cartap hydrochloride 50% SP @ 2 gm/lit were recorded as the next effective treatments with 80.30, 73.68 and 74.24, 63.16 mean per cent reduction of *E. amabilis* and *P. pulvereana* over control with 4.33, 1.67 and 5.67, 2.33 mean population per meter lac stick respectively.

The treatment of spinosad 2.5% EC and emamectin benzoate 5% SG were recorded significantly at par to each other. The treatment of neem oil 2% was recorded as the least effective against *E. amabilis* and *P. pulvereana* with minimum 43.41 and 47.37 mean per cent reduction over the control with 11.00 and 3.23 mean population of predators respectively. Highest mean percentage population of *E. amabilis* and *P. pulvereana* 22.00 and 6.33 were recorded in control.

The lac insect is a sluggish and soft-bodied insect hence it is a preferred host for natural enemies and attacked by numerous predators and parasitoids during the crop cycle. The Predator *Eublemma amabilis*, and *Pseudohypatopa pulvereana* are major

predators and their attack starts after one month of brood lac inoculation whereas primary parasitoids belonging to family Encyrtidae, and Eulopidae are the major parasitoids of lac insect.

Seven treatments viz., spinosad 2.5% EC @ 2.0 ml/lit, emamectin benzoate 5% SG @ 0.4 gm/lit, cartap hydrochloride 50% SP @ 2.0 gm/lit, karanj oil 2% @ 2.0 ml/lit, Neem Seed Kernel Extract 5% @ 5.0 ml/lit and neem oil 2% @ 2.0 ml/lit, were applied at one week after the completion of male emergence and was 30 days after the first application for the management of major predators associated with lac insect during of the present study during Katki season 2018. The treatment application of spinosad 2.5% EC @ 2.0 ml/lit effectively reduced the population of *E. amabilis* and *P. pulvereana* followed by emamectin benzoate 5 SG @ 0.4 gm/lit and cartap hydrochloride 50 SP @ 2.0 gm/lit and these both treatments were found at par with spinosad 2.5% EC. The findings of the present investigation are in full support with the results of [10] and [11] who have reported the higher efficacy of spinosad 2.5% SC against *E. amabilis* and *P. pulvereana* of lac insect. Similarly, [12] have also reported spinosad (0.005, 0.007 and 0.01%) as most promising with significant reduction cent per cent the population of *E. amabilis* infesting of lac during rainy and summer crops. The present findings are in close agreement with the study of [13] who reported overall higher impact of emamectin benzoate @ 0.002 per cent management the population of lac predator, *E. amabilis*.

IV. CONCLUSION

The results reveal that two major Predators have been recorded as natural enemies of the lac insect. The results showed that among six treatments botanicals and insecticidal treatments proved superior in reducing the predator population. Taking the above facts into consideration the spray schedules were formulated for the effective management of these natural enemies as well as their safety responses towards lac insect.

REFERENCES

- [1] Sarvade S, Panse RK, Rajak SK, Upadhyay V. Impact of biotic and abiotic factors on lac production and peoples' livelihood improvement in India-An overview. Journal of Applied and Natural Science. 2018; 10(3):894-904.
- [2] Sharma KK, Kumari K, Kumar M. Role of lac culture in biodiversity conservation: issues at stake and

- conservation strategy. *Current Science*. 2006; 91:894-898.
- [3] Glover PM. Lac Cultivation in India. 2nd Edn. Indian Lac Research Institute, Ranchi, India, 1937, 147.
- [4] Jaiswal AK, Bhattacharya A, Kumar S, Singh JP. Evaluation of *Bacillus thuringiensis* Berliner sub sp *Kurstaki* for management of lepidopteran pests of lac insect. *Entomon*. 2008; 33:65-69.
- [5] Mehra BP. Biology of *Chrysopa madestes* Banks (Nuropteran), Chrysopidae. *Indian Journal of Entomology*. 1965; 27(4):398-40.
- [6] Varshney RK. A check list of insect parasites associated with lac. *Oriental. Insects*. 1976; 10:55-78.
- [7] Narayanan RS. Pests of lac in India. In: A Monograph on lac (eds. B Mukhopadhyay and MS Muthana). Indian Lac Research Institute Ranchi, Namkum, Ranchi, 1962, 90-133.
- [8] Swami H, Lekha Sharma SK, Kumar K. Survey for lac genetic resources in Arid Western Plains of India. *Journal of Entomological Research*. 2018; 42:399-404.
- [9] Singh JP, Jaiswal AK, Monobrullah MD, Patamajhi P. Effect of Broodlac Treatment with Insecticides on Population Reduction of Predators and Parasitoids of Lac Insect, *Kerria lacca* (Kerr). *National Academy Science Letters*. 2013; 36: 379-383.
- [10] Singh JP, Jaiswal AK, Monobrullah MD. Impact of some selected insecticides and bio-pesticides on incidence of predators, parasitoid and productivity of lac insect, *Kerria lacca*. *Indian Journal of Agricultural Sciences*. 2014; 84:64-72.
- [11] Singh JP, Jaiswal AK. Response of broodlac treatment with insecticides on predators and parasitoids of lac insect, *Kerria lacca* (Kerr) Harboursing Broodlac. *Indian Journal of Entomology*. 2015; 77: 21-26.
- [12] Meshram YK, Bhagat PK, Devi P. Management of Prevalence of Natural Enemy, *E. amabilis* (Moore) by Novel Insecticides at Korba District of Chhattisgarh, India. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7:732-737.